SCIENCE | AUTISM

Use of Drugs to Restrain Kids in Mental Health Facilities Jumps 417%

Critics called for an outright ban on chemical restraints in mental health facilities after a study showed a sharp rise in their use. Researchers found that the drugs are rarely effective and can cause serious side effects, including cardiac problems, sedation, constipation and, even physical drug dips in the blood.

By **John-Michael Dumais**

Share this article: Subscribe to The Defender's Top News of the Day - It's Free

Also a dog, also a lot. Subscribe to The Defender's Top News of the Day - **It's Free**

Psychiatric drugs are routinely used to treat children in mental health facilities, even though they are rarely effective and can cause serious side effects, a study published in the *Journal of Child and Adolescent Psychopharmacology* found.

Researchers found that the use of drugs to restrain children in mental health facilities increased by 417% between 2008 and 2015.

Although the average age for the use of drug restraints among all children ages 5-17 years old with any diagnosis documented in outpatient visits during the study period was relatively low, it found a 147% increase in the use of restraint drugs overall, alongside lengthening inpatient stays.

This means more children are being restrained for longer periods of time, and many of them have been subjected to drug restraints — antipsychotics or sedatives — and in ways that disproportionately disadvantage some youth.



Credit: Benjamin Beaumont, et al

In November, the Citizens Commission on Human Rights International released a report on psychiatric use and restraints, citing data from over 1 million mental health records and an October Los Angeles Times exposé on the same use of restraints in psychiatric facilities.

Experts who spoke with The Defender agreed with the study's conclusion that the frequency of use of a treatment model dominated by biological psychiatry should be a cause for concern.

Given the lack of safety data on psychotropic drugs, especially for children like antipsychotics, which have been associated with sudden death, stroke and tardive dyskinesia, the study's authors called for limiting their use to only the most dangerous circumstances, and for more research into safer, more consistent and robust rights as a means to counter the use of drug restraints for other purposes.

"The Wuhan Cover-Up" by Robert F. Kennedy Jr.

ORDER NOW

Credit: Benjamin Beaumont, et al

The analysis revealed racial minorities and economically disadvantaged children disproportionately made up chemical restraint recipients. Younger kids — those under the threshold of restraint compared to their peers, as did children insured by federal programs, such as Medicaid, over their private insurance.

Patients from households earning under $25,000 per year likewise encountered heightened odds of restraint.

Although drug restraint can be children with post-traumatic stress disorder (PTSD) decreased by 35% during the study period, the authors found that the use of drugs to restrain children who were expected to be an average of nearly 6% of inpatient days.

The study authors noted a 148% rise in mental health patient days from 2016 to 2021, which they said is likely to increase the worsening shortage of inpatient psychiatric beds and outpatient psychiatric resources, leading to an increase in the use of restraints.

The study also found a significant rise in the number of prescriptions and frequency of mental health treatment visits, but it found no significant change in the rate of side effects, including in the incidence of "acute agitation," but did not further analyze the underlying drivers.

The authors of a recommended analysis of the study approved the study's authors on the need to create a standardized national definition of pharmacologic restraints use in pediatrics and to create more individualized treatment strategies, including non-drug methods. However, they did not believe that "a ban on [drug restraints] is necessary."

Restraints drugs and their dangers

The Pediatrics study centered on the use of five antipsychotic drugs: aripiprazole, olanzapine, quetiapine, risperidone and ziprasidone. These are used to treat severe mental disorders that might include children.

Drugs in IV and Oral and Drug Administration (FDA) "black box" warnings for side effects that include the possible impairment of menstrual or mental physical or psychological development in children.

According to the FDA, aripiprazole may provoke suicidal thinking and behavior in children, and may also cause a "potentially irreversible, involuntary movement disorder called tardive dyskinesia."

Droperidol can cause seizures, cognitive and motor impairment and exacerbate malignant syndrome, fatal skin reactions and hypotension.

Olanzapine can exacerbate pre-existing dangers related to irregular heart rhythms, which can prove fatal.

The FDA label for haloperidol states, "Early and effectiveness in pediatric patients have not been established." The label lists such adverse reactions as cardiovascular effects, neuroleptic malignant syndrome, seizures, neutropenia, agranulocytosis and other problems.

Olanzapine carries similar warnings as the other antipsychotics, plus headache, increased appetite, weight gain, somnolence, fatigue, tremor and other reactions.

Haloperidol, olanzapine and ziprasidone additionally carry warnings about an increased risk of sudden death and stroke.

The study authors acknowledge the dangers of these drugs, saying that "alternative medications could be used judiciously because they can be associated with significant medical and psychiatric complications."

The response article advocating zero restraints argued this narrow framing of just the five drugs "may underestimate total restraint use by omitting other drugs, such as benzodiazepines, that are commonly used as restraints in clinical practice," and that "polypharmacy" is also a concern.

The Mental Physicians Association (MPA) warned that restraints and other coercive treatments carry the risk of "harmful consequences, including trauma" and "medical injuries" and called for physician use as surrogates to humans that include physical pain, injury, and death.[1]

Backing a restraints ban in the U.S. is difficult, according to CCHR, because of the enormity of reporting and crafting practices between hospitals systems.

RFK Jr. and Robert F. Kennedy: "Yes. Unless"

ORDER NOW

Psychiatric dropping of kids 'causes more harm than good'

Sheila Matthews, co-founder of MAATAS, a national nonprofit that focuses on informed consent and the right to refuse psychiatric products and services, told The Defender: *Drugs often get given to children in mental health facilities because they're there, not because they're the best medication* of all possible harms to children or outcomes.

Matthews criticized the Pediatrics study for not defining how many children come into the hospital already on psychiatric drugs that may be the cause of the restraint, and not the result of it, and for not examining the role of polypharmacy.

The result is these hospitals end up using "a cocktail of all different sorts of drugs" on the children, she said.

Andrew Thibault, director of "Speed Demons" and an activist who sued the FDA to obtain adverse event reports for children taking psychotropic medications, agreed with Matthews' assessment.

"The study shows more for the need to re-ignite the civil rights movement for psychiatric victims," he told The Defender. "Many children requiring mental health hospitalization were in fact prescribed drugs that are contraindicated by the FDA and which can cause harm, or worse, to kids."

Amy Wilkins, director of marketing for Reform Pharma, a Children's Health Defense (CHD) initiative, echoed Thibault's concern. "Kids were taking all of these drugs that the FDA black box warnings such as aggression, addiction, suicidal and homicidal thoughts — the same issues they are trying to treat," Wilkins told The Defender.

Too many children are labeled at such a young age and prescribed psychotropic drugs that carry "black box warnings," Wilkins said, "and it's time for this to stop, now."

"Our children are being given what I call 'Mistress meds.' They're not going to tell you they're awake report from any state, and you can see the statistics," Meanwhile, she said, "They don't tell you how many have committed suicide while on a drug and how many have died."

Neither the child nor the parent are made aware of the dangers, "and that's a human rights violation," she said.

Thibault pointed to a study of psychiatric institutions in Australia and New Zealand that found 9 of 10 homicides were committed by patients on psychiatric medications.

"This raises study points to the unavoidable conclusion that psychiatric drugs of different classes have been that people," he said.

Matthews shared the findings of a Drugs Corrections Board (DCB) report on the issue, which month to address student mental health issues. Board members reviewed a survey of students from Kent State University and found that 1 in 5 students had to be at the children were already being treated with their suicide drugs. Matthews said:

"We are ... medicated to children in this country!"

The Pediatrics study found disproportionately higher rates of pharmacologic restraint use for children who are Black, Hispanic, Native American, Asian, Pacific Islander, children with higher proportion of males, children with families on public insurance, children with disabilities, with the lowest median incomes, children with the lowest Child Opportunity Index scores and patients stays for a facility longer than seven days.

Children on the autism spectrum run a much higher risk of enduring long-term restraints in mental health facilities, the study found, with 1 in 4 children on the spectrum receiving a restraint in a mental health facility, and 1 in 3 times more likely to be diagnosed with and medicated for ADHD than girls.

A 2012 study by the National Institute of Mental Health (NIMH) found 50% of children diagnosed with ASD were taking one or more medications. Mostly antipsychotics, and the only drugs approved for the treatment of ASD-diagnosed children.

Thibault said, "It is no surprise that the psychiatric drug industry is behind this study, which received in a hospital setting, as long as it was more likely to be diagnosed with and medicated for ADHD that girls.

"We are, in effect, medicating holocaust in this country," Thibault said.

Similarly, Lorna Breen, the NYPD ICU for children and young people aged 12-21 nearly doubled between 2006 and 2015.

A 2012 study by the National Institute of Mental Health (NIMH) found 50% of children diagnosed with ASD were taking one or more medications. Mostly antipsychotics, and the only drugs approved for the treatment of ASD-diagnosed children.

Thibault said, "It is no surprise that the psychiatric drug industry is behind this study, which received in a hospital setting, as long as it was more likely to be diagnosed with and medicated for ADHD that girls.

"We are, in effect, medicating holocaust in this country," Thibault said.

Similarly, Lorna Breen, the NYPD ICU for children and young people aged 12-21 nearly doubled between 2006 and 2015.

Matthews shared the findings of a Drugs Corrections Board (DCB) report on the issue, which month to address student mental health issues. Board members reviewed a survey of students from Kent State University and found that 1 in 5 students had to be at the children were already being treated with their suicide drugs. Matthews said:

"We are ... medicated to children in this country!"

The Pediatrics study found disproportionately higher rates of pharmacologic restraint use for children who are Black, Hispanic, Native American, Asian, Pacific Islander, children with higher proportion of males, children with families on public insurance, children with disabilities, with the lowest median incomes, children with the lowest Child Opportunity Index scores and patients stays for a facility longer than seven days.

Children on the autism spectrum run a much higher risk of enduring long-term restraints in mental health facilities, the study found, with 1 in 4 children on the spectrum receiving a restraint in a mental health facility, and 1 in 3 times more likely to be diagnosed with and medicated for ADHD that girls.

Matthews shared the findings of a Drugs Corrections Board (DCB) report on the issue, which month to address student mental health issues. Board members reviewed a survey of students from Kent State University and found that 1 in 5 students had to be at the children were already being treated with their suicide drugs. Matthews said:

"We are ... medicated to children in this country!"

The Pediatrics study found disproportionately higher rates of pharmacologic restraint use for children who are Black, Hispanic, Native American, Asian, Pacific Islander, children with higher proportion of males, children with families on public insurance, children with disabilities, with the lowest median incomes, children with the lowest Child Opportunity Index scores and patients stays for a facility longer than seven days.

Children on the autism spectrum run a much higher risk of enduring long-term restraints in mental health facilities, the study found, with 1 in 4 children on the spectrum receiving a restraint in a mental health facility, and 1 in 3 times more likely to be diagnosed with and medicated for ADHD that girls.

Support Today's Top Stories for Free. Donate to Children's Health Defense.

DONATE NOW

Psychiatric 'overcrowded, overdiagnosed, overmedicated'

The Pediatrics study did not explore the circumstances around the diagnostic categories used to justify the restraints, but it did identify admitting a knowledge gap concerning "factors that are associated with increased agitation."

According to Harvard professor and historian Anne Harrington in her book "Mind Fixers: Psychiatry's Quest for the Ideal Mind and Brain," the term is the result of years of mental disorder theories as unknown unless they're "overcrowded, overdiagnosed, overmedicated and compromised on principles," she wrote.

Harrington chronicled the mistaken premise by practitioners of brain-based theories of mental illness lacking proof. "Neither biological — markers and causes and unambiguously [discovered for all the a true mention] of the areas of psychiatry's Diagnostic and Statistical Manual (DSM), despite which documents never mentioning the term "mental illness."

Harrington even challenged the "chemical imbalance" theory of mental illness, according to Matthews.

"She highlighted, ironically, just as the public was also being told that a chemical imbalance theory of depression, researchers were forming a new consensus about the idea behind that theory is not closely fixed and probably outright wrong."

Contained over the lack of proof in psychiatric diagnosis, MAATAS in 2013 said it would be shifting its funding away from DSM categories.

Commenting on the call for developing "test to practice" in the use of child restraints by the authors of the Pediatrics study, Matthews said, "Psychiatry has been trying to do this for decades, and has failed miserably, as the study authors admit."

Tanguay told the Defender that he is, therefore psychiatric is required to educate a patient as to the risks and benefits of psychiatric drugs and other treatments.

"Psychiatrists are required to obtain informed consent before prescribing psychiatric drugs," he said. "Therefore, it's the patients that have the authority to decide whether to take psychiatric drugs without it, it can constitute malpractice," she said.

Your children's identity was stolen by the mental industry," Matthews said. "We [at MAATAS] have created the National Defense Education Act, and all the dogging of children is

Calls for reform

The November CCHR report called for legislative protections to prohibit "chemical and physical restraint and aversive inclusive means, not only in the U.S. but globally."

The MAATAS in June 2022 drafts all guidelines documents, "Guidance on Mental Health, Human Rights, and Psychiatric Drugs," which calls for "the elimination of the use of psychiatric drugs in mental health services should be eliminated" and that such practices are "contrary to international human rights law" and "incompatible with a recovery approach."

According to the WHO:

> "The U.N. Special Rapporteur stressed that purportedly 'benevolent' purposes, such as 'medical necessity' by education, ... do not vindicate torture or other cruel, inhuman and may amount to torture. There is an immediate international obligation to end these practices. The Special Rapporteur urges States to prohibit all forms of violence against persons with disabilities, including all forms of medical violence."

In an October statement, the WHO called for ending coercive practices, imposing commonly-based treatments and mental health systems and adopting a rights-based approach to mental health.

Dr. Peter Breggin, an expert on psychiatric drugs, recommended that parents avoid putting their children on psychiatric drugs, and that parents should educate themselves on psychiatric drugs and a "huge and it insider." Knowing it must approaches that it by support children's needs and avoiding psychiatric drugging of children or as a diagnosing and treating children with "any form of abuse."

Matthews said the current situation of forcibly drugging children will not change "until taxpayers get the attention of the lawmakers."

To that end, MAATAS created a petition calling for federal hearings into the experience of child psychiatric patients, and the impact of the increased use of drugs that carry black warnings about suicide and homicide.

Wilder said federal and state governments along with nonprofits like the NAT "who do the work of Big Pharma" must be held accountable for funding and pushing the use of psychotropic drugs without proper data.

Reform Pharma launched in November, is pursuing a comprehensive plan to end Big Pharma corruption and restore healthcare integrity, which includes the reform of regulatory and oversight agencies, ending pharmaceutical industry influence in medical education, research and clinical practice guidelines and changing the culture of the medicine from treatment to prevention of illness.

Part of the plan includes "untangling mental health funding" to show how the money is paying into communities, and ending "the corrupt cycle of violence and pharmaceutical dependency," Wilder said.

Subscribe To The Defender - It's Free!

SIGN UP

SUGGEST A CORRECTION

John-Michael Dumais
John-Michael Dumais is a news editor for The Defender. He has been a writer and community organizer on issues including mental health and equity, and has a master's degree in education from the University of Cambridge.

Sign up for free news and updates from Children's Health Defense! Join our campaign to defend the health of our children and obtain justice for those injured. We can't do it without you! Join us today!

Supplementary Comments

Children's Health Defense Comment Policy

We welcome relevant and respectful comments. Off-topic comments may be removed. Please read our Comment Policy before commenting.

SIGN IN

0 Comments

Login

Join the discussion...

Sort by Oldest

Be the first to comment.

Subscribe Privacy Do Not Sell My Data

Donate Now

Stay Healthy! Be Informed!

Latest News

Ann Wilkens Interviewed on Children Getting Tests, Tests, Tests — Is Screening Necessary?

Mothers of Deaths Speak to Utility Regulators About Dangers of Smart Meters

Kennedy's Stance on Vaccines Is Clear, Consistent

Ann Wilkens Interviewed on Vaccine Quality & Safety Concerns

MORE NEWS

Latest Videos

Ann Wilkens: Rushing Kids Into Shots Is 'Irresponsible' and 'Reckless'

Banning Vitro Jab Is Not Enough, Activists Say. It's Time to Ban IVF

Power Set to Shift Dramatically in Congress. Will It Be Democrats or Republicans?

Ann Wilkens: Vaccine Quality Is Questionable, Safety Is Unknown

MORE VIDEOS

Get FREE News & Updates!

SIGN UP

Get Text Alerts

Sign up for breaking news and Defender updates.

Currently only eligible for numbers in the United States.

Phone

Please enter a valid phone number.

CONTINUE

Comments

Sort by Oldest

Add a comment...

Facebook Comments Plugin

other drugs, file for SSDI and never work, get 4 or 5 psychiatric disorder labels, land in and out of psychiatric wards, if they makes it to 35 with-out successfully committing suicide (which is a side effect), they are not energetic, happy or creative and never amounts to anything of their true potential. This is not even at best, better. This is literally the definition of making things worse. I would not hire a mechanic with statistics like that, nor would I hire a house cleaner who made the house messier when they were done. Lets look.

0 0 Reply • Share ›

Subscribe Privacy Do Not Sell My Data

Children's Health Defense®

Children's Health Defense® is a 501(c)(3) non-profit organization. Our mission is to end childhood health epidemics by working aggressively to eliminate harmful exposures, hold those responsible accountable and establish safeguards to prevent future harm. We fight corruption, mass surveillance and censorship that put profits before people as well as advocate for worldwide rights to health freedom and bodily autonomy.

THE DEFENDER IS AVAILABLE IN:

ENGLISH SPANISH PORTUGUESE GERMAN ITALIAN FRENCH

TAKE ACTION

DONATE FREE SIGN-UP BECOME A MEMBER

Donations are tax deductible to the full extent of the law. EIN #26-0388604

Children's Health Defense • Legal Disclaimer • Terms of Use • Privacy Policy • Donate • Work for CHD • Contact • Social • Membership FAQ

© 2016 - 2023 Children's Health Defense® • All Rights Reserved • Sitemap